



UPPSALA
UNIVERSITET

Molecular Biotechnology Programme

Uppsala University School of Engineering

UPTEC X 07 056		Date of issue 2008-01	
Author Niclas Rollborn			
Title (English) Separation and determination of prostate-specific antigen isoforms in urine			
Title (Swedish)			
Abstract Prostate-specific antigen (PSA) in urine from healthy participants (ages between 24 and 64 years old) and from patients with malign prostate cancer was separated in different isoforms by using two chromatography technologies and an ultra sensitive immunochromatographic test (IKR). The results are showing different isoforms with different amount of sialic acid in the antennary complex, which can have mono-, di- and multi-antennary structure. Other results are that the normal range of PSA in urine is somewhere between 100 and 1000 µg PSA/L in urine and that there is a leakage of PSA to the urethra between the urine rounds but also that the prostate has a continuous secretion of PSA during an urine round.			
Keywords Prostate-specific antigen (PSA), isoform, urine, human, Lectin affinity chromatography, Concavalin A, Ion exchange chromatography (IEC), Immunochromatographic test (IKR).			
Supervisors Dr. Maria Lönnberg Uppsala University Dept. of Physical and Analytical Chemistry, Surface Biotechnology			
Scientific reviewer Adj. Prof. Jan Carlsson Uppsala University Dept. of Physical and Analytical Chemistry, Surface Biotechnology			
Project name		Sponsors	
Language English		Security Secret until 2010-11-01	
ISSN 1401-2138		Classification	
Supplementary bibliographical information		Pages 31	
Biology Education Centre Box 592 S-75124 Uppsala		Biomedical Center Tel +46 (0)18 4710000	
		Husargatan 3 Uppsala Fax +46 (0)18 555217	